

*Gastrointestinal Associates of Rockland*  
500 New Hempstead Rd New City NY 10956  
845-362-3200  
Fax: 845-290-8180

**MOVI PREP/DULCOLAX LAXATIVES  
PREPARATION FOR COLONOSCOPY**

*\*Unless directed by the physician, if you take heart or blood pressure medication, you may take with only a sip of water at your normally scheduled time as well as take your inhaler at your normally scheduled time. \*If you are taking Warfarin, Coumadin, Plavix, clopidogrel bisulfate, or diabetic medication, discuss this with the physician prior to your scheduled procedure date.*

- **STOP PRADAXA FIVE (5) DAYS PRIOR TO PROCEDURE.**
- **STOP EFFIENT SEVEN (7) DAYS PRIOR TO PROCEDURE.**
- No aspirin or aspirin containing products, Alka-seltzer, anti-arthritis or anti-inflammatory medications for seven (7) days prior to the scheduled procedure date. Tylenol is fine.
- No Metamucil, Fibercon, Citrucel or iron supplements five (5) days before your procedure.
- No vegetables for two (2) days before the procedure.

**PURCHASE THE FOLLOWING:**

- 1 BOX Dulcolax Laxatives (over the counter-NO rx needed)
- MoviPrep prescription from your local Pharmacy

*\*Please follow the preparation carefully and call the office if you have any questions.*

**24 HOURS BEFORE YOUR PROCEDURE**

Start a clear liquid diet - water, apple juice, coffee, tea (without milk or nondairy creamer), broth, jello, ices, sherbet, (nothing red or purple).

**STEP #1--- AT \_\_\_\_\_ THE DAY \_\_\_\_\_ YOUR PROCEDURE:**

Take all four (4) Dulcolax laxatives with water. Do not chew or crush! Continue clear liquid diet.

**STEP#2--- AT \_\_\_\_\_ THE DAY \_\_\_\_\_ YOUR PROCEDURE:**

Mix MoviPrep by emptying one (1) Pouch A and one (1) Pouch B into the disposable container. Add Lukewarm water or Crystal Light Lemonade to the top line of the container . Mix or shake to dissolve.

The MoviPrep container is divided by (four) 4 marks. Every 15 minutes, drink the solution down to the next mark (approx 8 oz), until the full liter is complete.

**STEP #3--- AT \_\_\_\_\_ THE DAY \_\_\_\_\_ YOUR PROCEDURE:  
REPEAT STEP #2**

**You may continue to drink clear liquids up to six hours prior to your procedure and then nothing by mouth until after your procedure time.**

\*You must have someone to drive you or accompany you home following the procedure due to the medication given at the time of the procedure. Please check with your insurance regarding pre-certification. **BRING YOUR MEDICATION LIST WITH YOU!!**